

High school geometry theorems

Hilbert's axiomatic system.
Formalized by: Sana Stojanovic Djurdjevic
Proved by: ArgoGeoChecker.

30.03.2018.

Theorem 1 (th_6_01.) *Assuming that $p \neq q$ and $q \neq r$ and $r \neq p$ and lines p and q intersect and lines q and r intersect and lines r and p intersect and $A \in p$ and $A \in q$ and $A \notin r$ and $B \notin p$ and $B \in q$ and $B \in r$ and $C \in p$ and $C \notin q$ and $C \in r$ it holds that $A \neq B$ and $B \neq C$ and $C \neq A$.*

Proof:

1. It holds that $A = B$ or $A \neq B$.
2. Assume that: $A = B$.
 3. From the facts $B \notin p$ and $A = B$ it holds that $A \notin p$.
 4. From the facts $A \notin p$ and $A \in p$ we get contradiction.
5. Assume that: $A \neq B$.
6. It holds that $A = C$ or $A \neq C$.
 7. Assume that: $A = C$.
 8. From the facts $C \notin q$ and $A = C$ it holds that $A \notin q$.
 9. From the facts $A \notin q$ and $A \in q$ we get contradiction.
 10. Assume that: $A \neq C$.
 11. From the fact $A \neq C$ it holds that $C \neq A$.
12. It holds that $B = C$ or $B \neq C$.
 13. Assume that: $B = C$.
 14. From the facts $C \in p$ and $B = C$ it holds that $B \in p$.
 15. From the facts $B \notin p$ and $B \in p$ we get contradiction.
 16. Assume that: $B \neq C$.
 17. The conclusion follows from the facts $A \neq B$ and $B \neq C$ and $C \neq A$.
18. The conjecture follows in all cases.
19. The conjecture follows in all cases.
20. The conjecture follows in all cases.

QED

Theorem 2 (th_6_02.) *Assuming that $p \neq q$ and $q \neq r$ and $r \neq p$ and lines p and q intersect and lines q and r intersect and lines r and p intersect and $A \in p$ and $A \in q$ and $A \notin r$ and $B \notin p$ and $B \in q$ and $B \in r$ and $C \in p$ and $C \notin q$ and $C \in r$ and $A \neq B$ and $B \neq C$ and $C \neq A$ it holds that $\neg \text{col}(A, B, C)$.*

Proof:

1. From the facts $A \neq B$ and $A \in q$ and $B \in q$ and $C \notin q$ it holds that $\neg col(A, B, C)$ (using *ax_D1a*).
2. The conclusion follows from the fact $\neg col(A, B, C)$.

QED

Theorem 3 (th_6_03.) *Assuming that $p \neq q$ and $q \neq r$ and $r \neq p$ and lines p and q intersect and lines q and r intersect and lines r and p intersect and $A \in p$ and $A \in q$ and $A \notin r$ and $B \notin p$ and $B \in q$ and $B \in r$ and $C \in p$ and $C \notin q$ and $C \in r$ and $A \neq B$ and $B \neq C$ and $C \neq A$ and $\neg col(A, B, C)$ there exist plane α , such that $A \in \alpha$ and $B \in \alpha$ and $C \in \alpha$.*

Proof:

1. From the fact $\neg col(A, B, C)$ there exist a plane α , where $A \in \alpha$ and $B \in \alpha$ and $C \in \alpha$ (using *ax_I4a*).
2. The conclusion follows from the facts $A \in \alpha$ and $B \in \alpha$ and $C \in \alpha$.

QED

Theorem 4 (th_6_04.) *Assuming that $p \neq q$ and $q \neq r$ and $r \neq p$ and lines p and q intersect and lines q and r intersect and lines r and p intersect and $A \in p$ and $A \in q$ and $A \notin r$ and $B \notin p$ and $B \in q$ and $B \in r$ and $C \in p$ and $C \notin q$ and $C \in r$ and $A \neq B$ and $B \neq C$ and $C \neq A$ and $\neg col(A, B, C)$ and $A \in \alpha$ and $B \in \alpha$ and $C \in \alpha$ it holds that $p \in \alpha$ and $q \in \alpha$ and $r \in \alpha$.*

Proof:

1. From the facts $A \neq B$ and $A \in q$ and $B \in q$ and $A \in \alpha$ and $B \in \alpha$ it holds that $q \in \alpha$ (using *ax_I6*).
2. From the fact $C \neq A$ it holds that $A \neq C$.
3. From the facts $A \neq C$ and $A \in p$ and $C \in p$ and $A \in \alpha$ and $C \in \alpha$ it holds that $p \in \alpha$ (using *ax_I6*).
4. From the facts $B \neq C$ and $B \in r$ and $C \in r$ and $B \in \alpha$ and $C \in \alpha$ it holds that $r \in \alpha$ (using *ax_I6*).
5. The conclusion follows from the facts $p \in \alpha$ and $q \in \alpha$ and $r \in \alpha$.

QED
